

Letters

RESEARCH LETTER

Novel Coronavirus Infection in Hospitalized Infants Under 1 Year of Age in China

Since December 8, 2019, an epidemic of coronavirus disease 2019 (COVID-19) has spread rapidly.¹ As of February 6, 2020, China reported 31 211 confirmed cases of COVID-19 and 637 fatalities.



Audio

Previous studies suggest that COVID-19 is more likely to infect older adult men, particularly those with chronic comorbidities.²⁻⁴ Few infections in children have been reported. We identified all infected infants in China and described demographic, epidemiologic, and clinical features.

Methods | For this retrospective study, we identified all hospitalized infants diagnosed with COVID-19 infection between December 8, 2019, and February 6, 2020, in China. The summary number and geographic location of new COVID-19 infections, released daily by the central government, were screened to identify infants (aged 28 days to 1 year). Demographic information, including age, sex, and geographic location, released anonymously by local health departments, were then retrieved and local hospitals and the Centers for Disease Control and Prevention were contacted for demographic data, family clustering (≥ 1 infected family member residing with the infant), linkage to Wuhan (residing in or visiting Wuhan or contact with visitors from Wuhan ≤ 2 weeks before the onset of infection), clinical features (symptoms at admission, dates of admission and diagnosis), treatment (intensive care unit or mechanical ventilation), prognosis (any severe complica-

tions, including death), and discharge date. Efforts were made to reach families of patients to confirm the information.

Nasopharyngeal swabs were collected during hospitalization. Real-time polymerase chain reaction testing was used to detect COVID-19 according to the recommended protocol. Infection was defined as at least 2 positive test results.

This study was approved by the institutional review board of Wuhan University School of Health Sciences. Informed consent was waived as part of a public health outbreak investigation.

Results | Nine infected infants were identified between December 8, 2019, and February 6, 2020 (Table). All patients were hospitalized. Seven patients were female. The youngest was aged 1 month and the oldest was 11 months. There were 2 patients from Beijing, 2 from Hainan, and 1 each from Guangdong, Anhui, Shanghai, Zhejiang, and Guizhou.

Four patients were reported to have fever, 2 had mild upper respiratory tract symptoms, 1 had no symptoms but tested positive for COVID-19 in a designated screening because of exposure to infected family members, and 2 had no information on symptoms available. The time between admission and diagnosis was 1 to 3 days.

Families of all 9 infants had at least 1 infected family member, with the infant's infection occurring after the family members' infection. Seven infants were reported to be either living in Wuhan or having family members who visited Wuhan, 1 had no direct linkage to Wuhan, and 1 had no information available. None of the 9 infants required intensive care or mechanical ventilation or had any severe complications.

Table. Characteristics of 9 Hospitalized Infants Infected With Coronavirus Disease 2019

Characteristic	Patient								
	1	2	3	4	5	6	7	8	9
Demographics									
Age	9 mo	11 mo	8 mo	10 mo	7 mo	1 mo 26 d	3 mo	3 mo 22 d	6 mo
Sex	Female	Female	Female	Male	Female	Female	Female	Female	Male
Symptoms at onset	Fever, peaking at 38.8 °C	Mild fever	None	NA	Fever	Runny nose; cough	Cough; sputum production	Fever	NA
Time between admission and diagnosis, d	1	1	3	3	1	1	1	1	2
Epidemiologic history									
No. of family members infected	2	1	5	1	2	2	2	1	1
Linkage to Wuhan	Yes	Yes	Yes	Yes	Yes	Yes	Yes	NA	No
Treatment									
Intensive unit care	No	No	No	No	No	No	No	No	No
Mechanical ventilation	No	No	No	No	No	No	No	No	No
Severe complications	No	No	No	No	No	No	No	No	No

Abbreviation: NA, not available.

Discussion | Based on the sources of data used in this study, 9 infants were infected with COVID-19 and were hospitalized in China between December 2019 and February 6, 2020. Given the number of infections reported, the number of infected infants identified was small. This may be due to a lower risk of exposure or incomplete identification due to mild or asymptomatic disease, rather than resistance to infection.^{1,5} However, this study showed that infants can be infected by COVID-19; the earlier stage of the COVID-19 epidemic primarily involved adults older than 15 years.²⁻⁴

Family clustering occurred for all infected infants. Infants who have infected family members should be monitored or evaluated and family clustering should be reported to ensure a timely diagnosis.

Seven of the 9 infant patients were female. Previous studies found higher percentages of infection in men than women.²⁻⁴ Whether female infants may be more susceptible to COVID-19 infection than male infants requires further study.

The study was limited by small sample size, inclusion only of infants who were hospitalized, and lack of inclusion of asymptomatic patients. Although a systematic and comprehensive search was made for relevant infections in infants, the epidemic is spreading rapidly and incomplete identification of cases is possible.

Because infants younger than 1 year cannot wear masks, they require specific protective measures. Adult caretakers should wear masks, wash hands before close contact with infants, and sterilize the infants' toys and tableware regularly.

Min Wei, MD
Jingping Yuan, MD, PhD
Yu Liu, PhD
Tao Fu, PhD
Xue Yu, MS
Zhi-Jiang Zhang, MD, PhD

Author Affiliations: Renmin Hospital of Wuhan University, Wuhan, China (Wei, Yuan, Fu); School of Management, Wuhan Institute of Technology, Wuhan,

China (Liu); School of Health Sciences, Wuhan University, Wuhan, China (Yu, Zhang).

Corresponding Author: Zhi-Jiang Zhang, MD, PhD, Department of Preventive Medicine, School of Health Sciences, Wuhan University, No. 185 Donghu Rd, Wuhan 430071, China (zhang22968@163.com).

Published Online: February 14, 2020. doi:[10.1001/jama.2020.2131](https://doi.org/10.1001/jama.2020.2131)

Author Contributions: Dr Zhang had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. Drs Wei, Yuan, and Liu contributed equally.

Concept and design: Zhang.

Acquisition, analysis, or interpretation of data: Wei, Yuan, Liu, Fu, Yu, Zhang.

Drafting of the manuscript: Wei, Zhang.

Critical revision of the manuscript for important intellectual content: Yuan, Liu, Fu, Yu, Zhang.

Statistical analysis: Liu, Yu, Zhang.

Obtained funding: Zhang.

Administrative, technical, or material support: Zhang.

Supervision: Zhang.

Conflict of Interest Disclosures: None reported.

Funding/Support: This study was funded by the National Natural Science Foundation of China (grant 81641123) and the Fundamental Research Funds for the Central Universities (grant 2042017kf0193).

Role of the Funder/Sponsor: The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Additional Contributions: We thank all patients involved in the study.

1. Paules CI, Marston HD, Fauci AS. Coronavirus infections—more than just the common cold [published online January 23, 2020]. *JAMA*. doi:[10.1001/jama.2020.0757](https://doi.org/10.1001/jama.2020.0757)

2. Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study [published online January 30, 2020]. *Lancet*. doi:[10.1016/S0140-6736\(20\)30211-7](https://doi.org/10.1016/S0140-6736(20)30211-7)

3. Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan [published online February 7, 2020]. *JAMA*. doi:[10.1001/jama.2020.1585](https://doi.org/10.1001/jama.2020.1585)

4. Li Q, Guan X, Wu P, et al. Early transmission dynamics in Wuhan, China, of novel coronavirus-infected pneumonia. *N Engl J Med* [published online January 29, 2020]. doi:[10.1056/NEJMoa2001316](https://doi.org/10.1056/NEJMoa2001316)

5. Del Rio C, Malani PN. 2019 Novel coronavirus—important information for clinicians [published online February 5, 2020]. *JAMA*. doi:[10.1001/jama.2020.1490](https://doi.org/10.1001/jama.2020.1490)